

Neutrino Mass Matrix in Terms of Up-Quark Masses

Masako BANDO¹ and Midori OBARA²

¹ *Aichi University, Aichi 470-0296, Japan*

² *Graduate School of Humanities and Sciences,
Ochanomizu University, Tokyo 112-8610, Japan*

Abstract

We demonstrate that under the "symmetric zero texture" with minimal Majorana mass matrix, neutrino masses and mixing angles are expressed in terms of up-quark masses, m_t, m_c, m_u . This provides interesting relations among neutrino mixing angles and up-type quark masses. Especially we predict $|U_{e3}| \leq 0.11$ even if we include the small mixing effects coming from charged lepton side. Also absolute masses of three neutrinos are predicted almost uniquely. This is quite in contrast to the case where bi-large mixings come from the charged lepton sector with non-symmetric charged lepton mass matrix.

¹E-mail address: bando@aichi-u.ac.jp

²E-mail address: midori@hep.phys.ocha.ac.jp

1 Introduction

Recent results from KamLAND [1] have established the Large Mixing Angle (LMA) solution [2]. Combined with the observations by Super-Kamiokande [3, 4] and SNO [5], this confirms that V_{MNS} has two large mixing angles [6, 7, 8]:

$$\sin^2 2\theta_{23} > 0.83 \text{ (99\% C.L.)}, \quad \tan^2 \theta_{12} = 0.86 \leq \sin^2 2\theta_{12} \leq 1, \quad (1.1)$$

with the mass squared differences

$$\Delta m_{32}^2 \sim 2.5 \times 10^{-3} \text{ eV}^2, \quad \Delta m_{21}^2 \sim 7 \times 10^{-5} \text{ eV}^2. \quad (1.2)$$

Now the question is why such a large difference can exist between the quark and lepton sectors. Within grand unified theories (GUTs), the Yukawa couplings of quarks and leptons to Higgs field are related each other. Can GUT predict two large mixing angles from some symmetry principle?

The neutrino mixing angles are expressed in terms of MNS matrix [9];

$$V_{MNS} = U_l^\dagger U_\nu, \quad (1.3)$$

where U_l and U_ν diagonalizes M_l and M_ν , respectively,

$$U_l^T M_l U_l = \text{diag}(m_e, m_\mu, m_\tau), \quad U_\nu^T M_\nu U_\nu = \text{diag}(m_{\nu_e}, m_{\nu_\mu}, m_{\nu_\tau}), \quad (1.4)$$

where M_ν is calculated from neutrino right-handed Majorana mass matrix (M_R) and Dirac mass matrix (M_{ν_D});

$$M_\nu = M_{\nu_D}^T M_R^{-1} M_{\nu_D}. \quad (1.5)$$

We call "up-road" ("down-road") option when such a large mixing angle comes from M_ν (M_l) side. In GUT framework, then, the up- (down-) quark mass matrix will play an important role. Here we assume the up-road option and make a semi-empirical analysis by adopting the so-called symmetric four zero texture. We shall show how we can reproduce bi-large mixing angles.

2 Symmetric Texture

First we make a comment on the so-called symmetric texture which has been extensively investigated by many authors [10]. In general symmetric texture, M_l is hierarchical mass matrix and never gives large mixing angles since down quark mass matrix, M_d is hierarchical. Thus symmetric textures dictate only up-road option. Can then hierarchical M_u be consistent with M_ν while M_{ν_D} is hierarchical? We shall show first that large mixing angles does not arise from M_ν

if we restrict ourselves to symmetric texture with $U(1)$ family structure. As we know well, the most popular mechanism which explains hierarchical structure of masses may be the so-called Froggatt-Nielsen mechanism [11] using anomalous $U(1)$ family quantum number.

Let us consider an example of 2-family model in which we have the same $U(1)$ charges to left-handed up-type fermions and the right-handed fermions, x_1, x_2 ($x_1 > x_2$), respectively. Then we get the form M_ν from Eq. (1.5) with general forms of $M_{\nu D}$ and M_R ,

$$M_{\nu D} \sim \begin{pmatrix} \lambda^{2x_1} & \lambda^{x_1+x_2} \\ \lambda^{x_1+x_2} & \lambda^{2x_2} \end{pmatrix}, \quad M_R^{-1} = \begin{pmatrix} a & c \\ c & b \end{pmatrix} \\ \rightarrow M_\nu \sim (a\lambda^{2x_1} + 2c\lambda^{x_1+x_2} + b\lambda^{2x_2}) \begin{pmatrix} \lambda^{2x_1} & \lambda^{x_1+x_2} \\ \lambda^{x_1+x_2} & \lambda^{2x_2} \end{pmatrix}. \quad (2.6)$$

This indicates that M_ν is always proportional to the hierarchical matrix, $M_{\nu D}$. Hence, unless the dominant terms are canceled accidentally by making fine tuning, it is impossible to get large mixing angles.

On the contrary, the above argument is no more valid if we choose the texture zero matrix in Eq. (2.6). Actually if we take zero texture, we obtain large mixing angle;

$$M_{\nu D} \sim \begin{pmatrix} \lambda^{2x_1} & 0 \\ 0 & \lambda^{2x_2} \end{pmatrix}, M_R \sim \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \rightarrow M_\nu \sim \lambda^{2x_i+2x_j} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (2.7)$$

The above example shows that M_ν is no more proportional to M_u (see more general discussion in a separate paper [12]).

3 GUT with Symmetric Texture

The informations of M_d and M_u are well established and popular. A simple example of quark mass matrices is symmetric "zero texture" [10]. Let us take the following forms of M_u and M_d which reproduce the observed quark and charged lepton masses as well as CKM mixing angles [13]. Then we get their relations in $SO(10)$ GUT;

$$M_U : \quad M_u \simeq \begin{pmatrix} 0 & \sqrt{m_u m_c} & 0 \\ \sqrt{m_u m_c} & m_c & \sqrt{m_u m_t} \\ 0 & \sqrt{m_u m_t} & m_t \end{pmatrix} \leftrightarrow M_{\nu D}, \quad (3.8)$$

Thus once we fix the representation of Higgs field in each matrix element, M_l and M_{ν_D} are uniquely determined from M_d and M_u , respectively.

$$M_D : \quad M_d \simeq \begin{pmatrix} 0 & \sqrt{m_d m_s} & 0 \\ \sqrt{m_d m_s} & m_s & \sqrt{m_d m_b} \\ 0 & \sqrt{m_d m_b} & m_b \end{pmatrix} \leftrightarrow M_l. \quad (3.9)$$

Here let us adopt a simple assumption that each elements of M_U and M_D is dominated by the contribution either from **10** or **126** of $SO(10)$ representation. There are 16 options for the Higgs configuration of M_U (see Table 1). We show that the following option of M_U , together with M_D (Georgi-Jarlskog type [14]) and the most economical form of M_R ;

$$\begin{aligned} M_U &= \begin{pmatrix} 0 & \mathbf{126} & 0 \\ \mathbf{126} & \mathbf{10} & \mathbf{10} \\ 0 & \mathbf{10} & \mathbf{126} \end{pmatrix}, \quad M_D = \begin{pmatrix} 0 & \mathbf{10} & 0 \\ \mathbf{10} & \mathbf{126} & \mathbf{10} \\ 0 & \mathbf{10} & \mathbf{10} \end{pmatrix}, \\ M_R &= \begin{pmatrix} 0 & rM_R & 0 \\ rM_R & 0 & 0 \\ 0 & 0 & M_R \end{pmatrix}. \end{aligned} \quad (3.10)$$

can reproduce all the masses and mixing angles of neutrinos consistently with present experiments.

4 Option S

Now each matrix element of M_{ν_D} is determined by multiplying an appropriate Clebsch-Gordan (CG) coefficient, 1 or -3 , and also M_ν are easily calculated from Eq. (1.5),

$$M_{\nu_D} = m_t \begin{pmatrix} 0 & a & 0 \\ a & b & c \\ 0 & c & d \end{pmatrix}, \quad M_\nu = \begin{pmatrix} 0 & \frac{a^2}{r} & 0 \\ \frac{a^2}{r} & 2\frac{ab}{r} + c^2 & c(\frac{a}{r} + 1) \\ 0 & c(\frac{a}{r} + 1) & d^2 \end{pmatrix} \frac{m_t^2}{m_R}. \quad (4.11)$$

Hence, because of the hierarchical structure of M_u , $a \ll b \sim c \ll 1$. In order to get large mixing angle θ_{23} , the first term of 2-3 element of M_ν in Eq. (4.11) should dominate and get the same order of magnitude as d^2 , namely, $r \sim \frac{ac}{d^2} \sim \sqrt{\frac{m_u^2 m_c}{m_t^3}} \sim 10^{-7}$. This indicates that the ratio of the the right-handed Majorana mass of 3rd generation to those of the first and second generations, is very small. Such kind of mechanism is well known as "seesaw enhancement" [15, 16, 17]. This tiny r is very welcome [15]; the right-handed Majorana mass of the third generation must become of order of GUT scale while those of the first and

Table 1: Classification of the up-type quark mass matrices.

Type	Texture1	Texture2	Texture3	Texture4
S	$\begin{pmatrix} 0 & \mathbf{126} & 0 \\ \mathbf{126} & \mathbf{10} & \mathbf{10} \\ 0 & \mathbf{10} & \mathbf{126} \end{pmatrix}$	$\begin{pmatrix} 0 & \mathbf{126} & 0 \\ \mathbf{126} & \mathbf{10} & \mathbf{10} \\ 0 & \mathbf{10} & \mathbf{10} \end{pmatrix}$		
A	$\begin{pmatrix} 0 & \mathbf{126} & 0 \\ \mathbf{126} & \mathbf{126} & \mathbf{126} \\ 0 & \mathbf{126} & \mathbf{126} \end{pmatrix}$	$\begin{pmatrix} 0 & \mathbf{126} & 0 \\ \mathbf{126} & \mathbf{126} & \mathbf{126} \\ 0 & \mathbf{126} & \mathbf{10} \end{pmatrix}$	$\begin{pmatrix} 0 & \mathbf{10} & 0 \\ \mathbf{10} & \mathbf{10} & \mathbf{10} \\ 0 & \mathbf{10} & \mathbf{126} \end{pmatrix}$	$\begin{pmatrix} 0 & \mathbf{10} & 0 \\ \mathbf{10} & \mathbf{10} & \mathbf{10} \\ 0 & \mathbf{10} & \mathbf{10} \end{pmatrix}$
B	$\begin{pmatrix} 0 & \mathbf{10} & 0 \\ \mathbf{10} & \mathbf{126} & \mathbf{126} \\ 0 & \mathbf{126} & \mathbf{126} \end{pmatrix}$	$\begin{pmatrix} 0 & \mathbf{10} & 0 \\ \mathbf{10} & \mathbf{126} & \mathbf{126} \\ 0 & \mathbf{126} & \mathbf{10} \end{pmatrix}$		
C	$\begin{pmatrix} 0 & \mathbf{126} & 0 \\ \mathbf{126} & \mathbf{10} & \mathbf{126} \\ 0 & \mathbf{126} & \mathbf{126} \end{pmatrix}$	$\begin{pmatrix} 0 & \mathbf{126} & 0 \\ \mathbf{126} & \mathbf{10} & \mathbf{126} \\ 0 & \mathbf{126} & \mathbf{10} \end{pmatrix}$	$\begin{pmatrix} 0 & \mathbf{10} & 0 \\ \mathbf{10} & \mathbf{10} & \mathbf{126} \\ 0 & \mathbf{126} & \mathbf{126} \end{pmatrix}$	$\begin{pmatrix} 0 & \mathbf{10} & 0 \\ \mathbf{10} & \mathbf{10} & \mathbf{126} \\ 0 & \mathbf{126} & \mathbf{10} \end{pmatrix}$
F	$\begin{pmatrix} 0 & \mathbf{126} & 0 \\ \mathbf{126} & \mathbf{126} & \mathbf{10} \\ 0 & \mathbf{10} & \mathbf{126} \end{pmatrix}$	$\begin{pmatrix} 0 & \mathbf{126} & 0 \\ \mathbf{126} & \mathbf{126} & \mathbf{10} \\ 0 & \mathbf{10} & \mathbf{10} \end{pmatrix}$	$\begin{pmatrix} 0 & \mathbf{10} & 0 \\ \mathbf{10} & \mathbf{126} & \mathbf{10} \\ 0 & \mathbf{10} & \mathbf{126} \end{pmatrix}$	$\begin{pmatrix} 0 & \mathbf{10} & 0 \\ \mathbf{10} & \mathbf{126} & \mathbf{10} \\ 0 & \mathbf{10} & \mathbf{10} \end{pmatrix}$

second generations are of order 10^8 GeV. This is quite favorable for the GUT scenario to reproduce the bottom-tau mass ratio. With such a tiny r , M_ν is approximately written as,

$$M_\nu = \begin{pmatrix} 0 & \frac{a^2}{r} & 0 \\ \frac{a^2}{r} & \frac{2ab}{r} & \frac{ac}{r} \\ 0 & \frac{ac}{r} & d^2 \end{pmatrix} \frac{m_t^2}{m_R} \equiv \begin{pmatrix} 0 & \beta & 0 \\ \beta & \alpha & h \\ 0 & h & 1 \end{pmatrix} \frac{d^2 m_t^2}{m_R}, \quad \begin{aligned} h &= ac/rd^2, \\ \alpha &= 2ab/rd^2, \\ \beta &= a^2/rd^2, \end{aligned} \quad (4.12)$$

where $\beta \ll \alpha$ and $h \sim \mathcal{O}(1)$.

First let us diagonalize the dominant term with respect to 2-3 submatrix of Eq. (4.12) with the rotation angle θ_{23} ,

$$\xrightarrow{\theta_{23}} \begin{pmatrix} 0 & \beta \cos \theta_{23} & \beta \sin \theta_{23} \\ \beta \cos \theta_{23} & \lambda_2 & 0 \\ \beta \sin \theta_{23} & 0 & \lambda_{\nu_3} \end{pmatrix}, \quad \tan^2 2\theta_{23} = \frac{4h^2}{(1-\alpha)^2}, \quad (4.13)$$

with their eigenvalues,

$$\lambda_{\nu_3} = \frac{\alpha + 1 + \sqrt{(\alpha - 1)^2 + 4h^2}}{2}, \quad \lambda_2 = \frac{\alpha + 1 - \sqrt{(\alpha - 1)^2 + 4h^2}}{2}. \quad (4.14)$$

Second step is to rotate with respect to 1 and 2 components of Eq. (4.13);

$$\begin{aligned} \xrightarrow{\theta_{12}} & \begin{pmatrix} \lambda_{\nu_1} & 0 & \beta \sin \theta_{23} \cos \theta_{12} \\ 0 & \lambda_{\nu_2} & \beta \sin \theta_{23} \sin \theta_{12} \\ \beta \sin \theta_{23} \cos \theta_{12} & \beta \sin \theta_{23} \sin \theta_{12} & \lambda_{\nu_3} \end{pmatrix}, \\ \tan^2 2\theta_{12} &= \left(\frac{2\beta \cos \theta_{23}}{\lambda_2} \right)^2, \end{aligned} \quad (4.15)$$

with eigenvalues,

$$\lambda_{\nu_2} = \frac{\lambda_2 + \sqrt{\lambda_2^2 + 4\beta^2 \cos^2 \theta_{23}}}{2}, \quad \lambda_{\nu_1} = \frac{\lambda_2 - \sqrt{\lambda_2^2 + 4\beta^2 \cos^2 \theta_{23}}}{2}. \quad (4.16)$$

Finally the neutrino masses are given as,

$$m_{\nu_3} \sim \lambda_{\nu_3} \frac{d^2 m_t^2}{m_R}, \quad m_{\nu_2} \sim \lambda_{\nu_2} \frac{d^2 m_t^2}{m_R}, \quad m_{\nu_1} \sim \lambda_{\nu_1} \frac{d^2 m_t^2}{m_R}. \quad (4.17)$$

5 Numerical Calculations

Using the up-quark masses at GUT scale within the error [18],

$$m_u = 1.04_{-0.20}^{+0.19} \text{ MeV}, \quad m_c = 302_{-27}^{+25} \text{ MeV}, \quad m_t = 129_{-40}^{+196} \text{ GeV}, \quad (5.18)$$

the parameter range of $\alpha = 2hb/c$ and $\beta = ha/c$ are estimated from the following values;

$$\frac{2b}{c} \rightarrow \frac{2m_c}{\sqrt{m_u m_t}} \sim 1.0 - 2.4, \quad \frac{a}{c} \rightarrow \sqrt{\frac{m_c}{m_t}} \sim 0.03 - 0.05. \quad (5.19)$$

In order to realize large mixing angle θ_{23} , the option in which α is close to 1 is a better choice. On the other hand, in order to realize large mixing angle θ_{12} , λ_2 must become at least of the same order as 2β , so the option in which β is relatively large would be a better choice. Thus the desired candidate for the options of Table 1 would be 1) *The Higgs representations coupled with 2-3 and 2-2 elements of M_U must be same.* 2) *The Higgs representation coupled with 1-2 elements of M_U must be as large as possible.* The option S may be the best candidates which satisfy the conditions (i) and (ii).

Leaving the detailed calculations to our full paper [12], we here show an example of the figures of our results in Fig. 1. The explicit forms of up-type mass matrix for the class S are seen in Eq. (3.9) with Eq. (3.10), which we expected in section 2. Those two types yield the same predictions except for




Figure 1: Calculated values of $\sin^2 2\theta_{23}$ versus h and $\tan^2 \theta_{12}$ versus h in the class S . The experimentally allowed regions are indicated by the horizontal lines.

the Majorana mass scale. The type S_1 requires $m_R \sim 2 \times 10^{15}$ GeV and in the type S_2 , we have $m_R \sim 10^{14}$ GeV, respectively. Thus more desirable one may be the type S_1 since it predicts more realistic bottom-tau ratio at low energy. Then the neutrino mass matrix is written as,

$$M_\nu \sim \begin{pmatrix} 0 & -3h\sqrt{\frac{m_c}{m_t}} & 0 \\ -3h\sqrt{\frac{m_c}{m_t}} & 2h\frac{m_c}{\sqrt{m_u m_t}} & h \\ 0 & h & 1 \end{pmatrix} \frac{9m_t^2}{m_R}. \quad (5.20)$$

From this, we obtain the following equations,

$$\tan^2 2\theta_{23} \simeq \frac{4h^2}{\left(1 - h\frac{2m_c}{\sqrt{m_u m_t}}\right)^2}, \quad \tan^2 2\theta_{12} \simeq \frac{144h^2 m_c \cos^2 \theta_{23}}{m_t \left(1 - 2h + h\frac{2m_c}{\sqrt{m_u m_t}}\right)^2}. \quad (5.21)$$

The neutrino masses are given by

$$m_{\nu_3} \simeq \lambda_{\nu_3} \cdot \frac{m_t^2}{m_R}, \quad m_{\nu_2} \simeq \lambda_{\nu_2} \cdot \frac{m_t^2}{m_R}, \quad m_{\nu_1} \simeq \lambda_{\nu_1} \cdot \frac{m_t^2}{m_R}, \quad (5.22)$$

where the RGE factor ($\sim 1/3$) has been taken account in estimating lepton masses at low energy scale. Since $\lambda_{\nu_2} \ll \lambda_{\nu_3} \sim \mathcal{O}(1)$, this indeed yields $m_R \sim 10^{15}$ GeV, as many people require. We here list a set of typical values of neutrino masses and mixings at $h = 0.9$, $m_t \simeq 260$ GeV;

$$\sin^2 2\theta_{23} \sim 0.98 - 1, \quad \tan^2 \theta_{12} \sim 0.29 - 0.46, \quad (5.23)$$

$$m_{\nu_3} \sim 0.053 - 0.059 \text{ eV}, \quad m_{\nu_2} \sim 0.003 - 0.008 \text{ eV}, \quad (5.24)$$

$$m_{\nu_1} \sim 0.0006 - 0.001 \text{ eV}. \quad (5.25)$$

Up to here we have estimated the contribution from M_ν side and this is fairly in a good approximation for estimating large mixing angles θ_{23} and θ_{12} . Also

neutrino masses are determined from M_ν only. However, in estimating $|U_{e3}|$, we need careful estimation, since we cannot neglect the additional contribution from the charged lepton side in Eq. (1.3). The contribution from M_ν is as follows,

$$\sin \theta_{13} \simeq \frac{6h}{1 + 2h + h \frac{2m_c}{\sqrt{m_u m_t}}} \sqrt{\frac{m_c}{m_t}} \sin \theta_{23} \cos \theta_{12}. \quad (5.26)$$

Within the allowed range of h , the calculated value of θ_{13} from M_ν is almost of the same order as the one from M_l ;

$$|\theta_{13}^\nu| \sim 0.037 - 0.038 \quad \leftrightarrow \quad \Delta\theta_{13}^l \sim \frac{\lambda}{3} \cdot U_{\mu 3} \sim 0.04, \quad (5.27)$$

where $\lambda \sim 0.2$ and the factor 3 comes from the Georgi-Jarlskog texture of Eq. (3.10). We have to combine those two contributions; unfortunately we do not yet have exact information of the relative phase. If the two terms act additively (negatively), we would have maximal (minimum) value. Still we can say that $|U_{e3}|$ becomes at most 0.11, which is within the experimental limit [19]. This would be one of the very important predictions of this model. In order to predict exact $|U_{e3}|$, the inclusion of CP phase of M_ν and M_l is important, which is our next task. In conclusion we have seen that the up-road option can reproduce the present neutrino experimental data very well. However also down-road option may be also worthwhile to be investigated [20], in which case the Nature may show "twisted family structure". On the contrary in the case of up-road option it requires "parallel family structure".

Acknowledgements

We would like to thank to M. Tanimoto, A. Sugamoto and T. Kugo whose stimulating discussion encouraged us very much. Also we are stimulated by the fruitful and instructive discussions at the Summer Institute 2002 held at Fuji-Yoshida and at the research meeting held in Nov. 2002 supported by the Grant-in Aid for Scientific Research No. 09640375. M. B. is supported in part by the Grant-in-Aid for Scientific Research No. 12640295 from Japan Society for the Promotion of Science, and Grants-in-Aid for Scientific Purposes (A) "Neutrinos" (Y. Suzuki) No. 12047225, from the Ministry of Education, Science, Sports and Culture, Japan.

References

- [1] KamLAND Collaboration, K. Eguchi *et al.*, Phys. Rev. Lett. **90** (2003) 021802.

- [2] P.C.de Holanda and A. Smirnov, hep-ph/0212270
- [3] Super-Kamiokande Collaboration, S. Fukuda *et al.*, Phys. Lett. **B433** (1998) 9; *ibid.* **436**, 33 (1998); *ibid.* **539**, 179 (2002).
- [4] Super-Kamiokande Collaboration, S. Fukuda *et al.*, Phys. Rev. Lett. **86** (2001) 5651; *ibid.* **86**, 5656 (2001).
- [5] SNO Collaboration, Q.R. Ahmad *et al.*, Phys. Rev. Lett. **89** (2002) 011301; *ibid.* **89**, 011302 (2002).
- [6] M. Maltoni, T. Schwetz and J.W.F. Valle, arXiv:hep-ph/0212129.
- [7] G.L. Fogli *et al.*, arXiv: hep-ph/0212127.
- [8] J.N. Bahcall, M.C. Gonzalez-Garcia, and C. Pena-Garay, arXiv:hep-ph/0212147.
- [9] Z. Maki, M. Nakagawa and S. Sakata, Prog. Theor. Phys. **28** (1962) 870.
- [10] P. Ramond, R.G. Roberts and G.G. Ross, Nucl. Phys. **B406** (1993) 19; J. Harvey, P. Ramond and D. Reiss, Phys. Lett. **B92** (1980) 309; Nucl. Phys. **B199** (1982) 223; S. Dimopoulos, L.J. Hall and S. Raby, Phys. Rev. Lett. **68** (1992) 1984; Phys. Rev. **D45** (1992) 4195; Y. Achiman and T. Greiner, Nucl. Phys. **B443** (1995) 3; D. Du and Z.Z. Xing, Phys. Rev. **D48** (1993) 2349; H. Fritzsch and Z.Z. Xing Phys. Lett. **B353** (1995) 114; K. Kang and S.K. Kang, Phys. Rev. **D56** (1997) 1511; J.L. Chkareuli and C.D. Froggatt, Phys. Lett. **B450** (1999) 158; P.S. Gill and M. Gupta, Phys. Rev. **D57** (1998) 3971; M. Randhawa, V. Bhatnagar, P.S. Gill and M. Gupta, Phys. Rev. **D60** (1999) 051301; S.K. Kang and C.S. Kim, Phys. Rev. **D63** (2001) 113010; M. Randhawa, G. Ahuja and M. Gupta, arXiv: hep-ph/0230109.
- [11] C. D. Frogatt and H. B. Nielsen, Nucl. Phys. **B147** (1979) 277.
- [12] M. Bando and M. Obara, arXiv: hep-ph/0212242.
- [13] H. Nishiura, K. Matsuda and T. Fukuyama, Phys. Rev. **D60** (1999) 013006. K. Matsuda, T. Fukuyama and H. Nishiura, Phys. Rev. **D61** (2000) 053001.
- [14] H. Georgi and C. Jarlskog, Phys. Lett. **B86** (1979) 297.
- [15] M. Bando, T. Kugo and K. Yoshioka, Phys. Rev. Lett. **80** (1998) 3004.
- [16] M. Tanimoto, Phys. Lett. **B345** (1995) 477.

- [17] A. Yu. Smirnov, Phys. Rep. **D48** (1993) 3264.
- [18] H. Fusaoka and Y. Koide, Phys. Rev. **D57** (1998) 3986.
- [19] M. Apollonio *et al.* (CHOOZ), Phys. Lett. **B420** (1998) 397.
- [20] T. Yanagida, Talk given at 18th Int. Conf. on Neutrino Physics and Astrophysics (NEUTRINO 98), Takayama, Japan, 4-9 Jun 1998; hep-ph/9809307; J. Sato and T. Yanagida, Phys. Lett. **B430** (1998) 127; Y. Nomura and T. Yanagida, Phys. Rev. **D59** (1998) 017303; H. Haba and H. Murayama, Phys. Rev. **D63** (2001) 053010; M. Bando and T. Kugo, Prog. Theor. Phys. **101** (1999) 1313; M. Bando, T. Kugo and K. Yoshioka, Phys. Lett. **B483** (2000) 163.